LEMTRADA® Referral Form



Fax completed form to:

PATIENT INFORMATION		
Patient Name:	Date of Birth: Referral Date:	
Address:	City/State/Zip:	
Home Phone:	Cell Phone: Work Phone:	
Secondary Contact:	Height: Weight: Male Female	
Patient Diagnosis & ICD)-10:	
Allergies:		
PROVIDER INFORMATION		
Physician Name:	Lic.#: DEA #:	
Practice Name:	NPI#:	
Address:	City/State/Zip:	
Office Contact:	Phone: Fax:	
Supervisory Physician (i		
	MS CLINICAL DETAILS	
Type of MS: Primary progressive multiple sclerosis (PPMS) OR Relapsing multiple sclerosis (RMS)		
Ambulation status:	Able to ambulate more than 5 meters Able to ambulate without aid or rest for at least 100 meters	
Relapse details: Two or more relapses within the previous two years One relapse within the previous year		
PLEASE ATTACH		
Patient demographi	ics & front/back copy of all insurance cards (prescription & medical) CBC with differential, Serum creatinine levels, urinalysis with cell counts, urine protein to cre	atinine ratio
Recent office visit notes, history & physical, lab & pertinent procedure results		
Current modification list & list of current modification triad and failed (with data)		
vaccine status (any vaccination) and documentation of any fecent vaccinations		
Line access docume	entation/verification if applicable Letter of medical necessity if drug dosing or indication is outside of FDA guidelines	
	NURSING & LAB ORDERS	
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion.		
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line		
	/M per nasal cannula as needed	
Lab Orders:	Lab Date & Frequency:	
SUPPLY ORDERS		
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary		
		REFILLS
PRODUCT	PRESCRIPTION INFORMATION	KEFILLS
Is this a first dose?	Yes No If No, when was last dose given? When is patient due for next dose?	
	Pre Meds: Hydroxyzine HCl 50mg po prior to start of infusion and every 6 hours prn #25	
	Acyclovir 200mg po BID for a minimum of 2 months or until CD4+ count is > or = to 200 cells per microliter, whichever occurs later #60 Refill: #1	
	Cetirizine 10mg po prior to Lemtrada infusion Ondansetron 4mg po prn #25	
Lemtrada	Promethazine 25mg po prn #25 Famotidine 20mg prior to start of alemtuzumab infusion	
	Acetaminophen 1000mg po prior to start of Lemtrada infusion and q6h prn Other:	
	Note – If needed, please send pain prescription to retail pharmacy	
	Pre Infusion: Solu-Medrol 1000mg IV infusion in 500mL of 0.9% NaCl over 1 hour prior to Lemtrada infusion on days 1, 2 and 3 only	
	Normal saline 0.9% 500ml IV prior to Lemtrada infusion on days 4 and 5	
	Initial Course: 12mg/day IV infusion over 4 hours for 5 consecutive days	
	Subsequent Course: 12mg/day IV infusion over 4 hours for 3 consecutive days *To start at least12 months after previous dose*	
	Post Meds: Normal saline 0.9% 500mL IV infusion over 1 hour post Lemtrada infusion	
	Ondansetron 4-8mg in 50-100mL 0.9% NaCl IV infusion over 15 minutes prn nausea	
	Epinephrine 1:1000 0.5-1mL IVP prn angioedema/hypotension/branchospasm/generalized urticaria	
/ SIDE EFFECT ORDERS	Ketorolac: 30mg IVP over 3-5 minute	
	Diphenhydramine 50mg in 100mL of 0.9% NaCl IV over approx 15 mins prn pruitis/rash	
071150		
OTHER	nd utilizing our services, you are authorizing EventusRx to serve as your prior authorization designated agent in dealing with medical and prescription insuranc	

Date

Prescriber's Signature Substitution Permitted Print Name

