

Soliris® Order Form



Fax Completed Form To:

Phone:

PATIENT INFORMATION			
Patient Name:		Date of Birth:	
Address:		Referral Date:	
Home Phone:		City/State/Zip:	
Cell Phone:		Work Phone:	
Secondary Contact:	Height:	Weight:	☐ Male ☐ Female
Patient Diagnosis & ICD-10:			
Allergies:			
PROVIDER INFORMATION			
Physician Name:		Lic.#:	
Practice Name:		DEA #:	
Address:		NPI#:	
Office Contact:		City/State/Zip:	
Phone:		Fax:	
Supervisory Physician (if applicable):			
PLEASE ATTACH			
☐ Patient demographics & front/back copy of all insurance cards (prescription & medical) ☐ Recent office visit notes, history & physical, lab & pertinent procedure results ☐ Current medication list & list of prior medications tried and failed (with dates) ☐ Line access documentation/verification if applicable		☐ Vaccine status (any vaccination) and documentation of any recent vaccinations ☐ Clinical documentation on any recent meningococcal infections ☐ Documentation of a meningococcal vaccination ☐ Letter of medical necessity if drug dosing or indication is outside of FDA guidelines	
NURSING & LAB ORDERS			
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - ☐ 10units/mL ---OR--- ☐ 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders:			
Lab Date & Frequency:			
PRESCRIPTION ORDERS			
Anaphylaxis Kit: ☐ Epinephrine 0.3mg IM as needed ☐ Solu-cortef 250mg-500mg IV infusion as needed ☐ Solu-Medrol 60mg - 125mg IV infusion as needed (Check all that apply) ☐ Diphenhydramine _____ mg IV infusion as needed ☐ NS Hydration 500 ml IV infusion over 30 minutes as needed ☐ Other			
Pre-Medications: ☐ Acetaminophen _____ mg PO _____ minutes prior to infusion ☐ Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion (Check all that apply) ☐ Diphenhydramine _____ mg ☐ PO ---OR--- ☐ IV infusion _____ minutes prior to infusion ☐ Other			
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			
PRODUCT	PRESCRIPTION INFORMATION		REFILLS
Is this a first dose? ☐ Yes ☐ No If No, when was last dose given? _____ When is patient due for next dose? _____			
Is the prescriber enrolled in the Soliris REMS program? ☐ Yes ☐ No			
☐ Soliris Induction (≥18 years of age)	☐ PNH ☐ 600 mg IV infusion via ☐ gravity ---OR--- ☐ pump every 7 days for 4 weeks over 35 minutes ☐ aHUS, gMG and NMOsD ☐ 900 mg IV infusion via ☐ gravity ---OR--- ☐ pump every 7 days for 4 weeks over 35 minutes		NONE
☐ Soliris Maintenance (≥18 years of age)	☐ PNH ☐ 900 mg IV infusion via ☐ gravity or ☐ pump every 2 weeks starting week 5 over 35 minutes ☐ aHUS, gMG and NMOsD ☐ 1,200 mg IV infusion via ☐ gravity or ☐ pump every 2 weeks starting week 5 over 35 minutes		_____
☐ Soliris Induction (<18 years of age)	aHUS ☐ For patients 5-10kg administer 300mg IV infusion via ☐ gravity ---OR--- ☐ pump once weekly X 1 dose over 1 to 4 hours ☐ For patients 10-20kg administer 600mg IV infusion via ☐ gravity ---OR--- ☐ pump once weekly X 1 dose over 1 to 4 hours ☐ For patients 20-30kg administer 600mg IV infusion via ☐ gravity ---OR--- ☐ pump once weekly X 2 doses over 1 to 4 hours ☐ For patients 30-40kg administer 600mg IV infusion via ☐ gravity ---OR--- ☐ pump once weekly X 2 doses over 1 to 4 hours ☐ For patients >40kg administer 900 mg IV infusion via ☐ gravity ---OR--- ☐ pump once weekly X 4 doses over 1 to 4 hours		NONE
☐ Soliris Maintenance (<18 years of age)	aHUS ☐ For patients 5-10kg administer 300 mg IV infusion via ☐ gravity ---OR--- ☐ pump starting at week 2 then 300mg every 3 weeks over 1 to 4 hours ☐ For patients 10-20kg administer 300 mg IV infusion via ☐ gravity ---OR--- ☐ pump starting at week 2 then 300mg every 2 weeks over 1 to 4 hours ☐ For patients 20-30kg administer 600 mg IV infusion via ☐ gravity ---OR--- ☐ pump starting at week 3 then 600mg every 2 weeks over 1 to 4 hours ☐ For patients 30-40kg administer 900mg IV infusion via ☐ gravity ---OR--- ☐ pump starting at week 3 then 900mg every 2 weeks over 1 to 4 hours ☐ For patients >40kg administer 1,200mg IV infusion via ☐ gravity ---OR--- ☐ pump starting at week 5 then 1,200mg every 2 weeks over 1 to 4 hours		_____
☐ OTHER			_____
By signing this form and utilizing our services, you are authorizing Amerita to assist with prior authorization requests acting as your pharmacy provider in dealing with medical and prescription insurance companies.			

Prescriber's Signature
Dispense as Written

Print Name

Date

Prescriber's Signature
Substitution Permitted

Print Name

Date

